

Physical Therapy Prescription – Meniscus Repair

Name:	Date of Surgery:
Procedure: R / L knee arthroscopy, meniscus re	epair
Frequency: 2-3 times per week for 6 weeks	
PHASE I (Weeks 0 – 6): Period of protection, decrease	edema, activate quadriceps
•	mbulation and sleeping tion and removed while sleeping d; no weight-bearing with knee flexion angles >90° mstring sets, heel slides, Gastroc/Soleus stretching, straight strength prevents extension lag
Phase II (Weeks 6 – 12)	
 Weightbearing: As tolerated, unassisted Hinged Knee Brace: Discontinue at 6 weeks Range of Motion: Full Therapeutic Exercises: Patellar mobs, quad/hacurls, toe raises, balance exercises, Gastroc/Soleo Begin use of the stationary bicycle Modalities: Per therapist, including electrical stire 	•
Phase III (Weeks 12 – 16)	
Range of Motion: Full, painless	

- - Therapeutic Exercises: Advance closed chain strengthening exercises and proprioception activities; focus on single-leg strengthening; begin elliptical
 - Straight ahead running permitted at 12 weeks
 - Swimming okay at 16 weeks
 - Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase IV (Weeks 16 - 24): Gradual return to athletic activity

- 16 weeks: begin jumping
- 20 weeks: advance to sprinting, backward running, cutting/pivoting/changing direction
- Consider functional sports assessment

Signature:	Date:
------------	-------