



Jill McAngus, MD

Distal Biceps Repair – Postoperative Instructions

Wound Care:

- After surgery, the wound is covered with gauze pads, web roll, posterior mold splint and ace wrap. These should be left in place for until first post-operative visit. If bright red blood soaks through the dressings, please call Dr. McAngus' office

Medications:

- Pain Control:
 - For most patients, a nerve block is provided by the anesthesia team before surgery to help with postoperative pain control – while every patient is different, this will typically wear off within 12-24 hours. Most patients will require some narcotic pain medication (i.e. vicodin, norco, percocet, or other codeine-derivative) for 1-2 days after surgery – please take as instructed.
 - It is important not to drink alcohol or drive while taking narcotic medication. If your pain is minimal, you may discontinue the use of narcotics. Ibuprofen 200-400 mg (i.e. Advil) can be taken as needed in between doses of narcotic pain medication for additional pain control.
- Constipation:
 - The use of narcotics can lead to constipation. Adequate hydration and over-the-counter stool softeners can minimize constipation problems.
- Normal medications:
 - Resume the day after surgery unless otherwise instructed.

Activity:

- ****No active triceps extension for 6 weeks****
- Unless otherwise instructed, you should begin simple hand and wrist exercises. You may use your arm to assist with eating and personal hygiene unless specifically instructed not to by Dr. McAngus.
- You may not bear-weight with your operative arm. Be sure to use and move your hand, wrist, and elbow in order to decrease swelling in your arm.
- While exercise is important, don't over-do it. Common sense is the rule. •
- Sling/Immobilizer:
 - Unless otherwise instructed, you should wear your postoperative sling/immobilizer at all times, including while sleeping. The sling can be taken off for showering; however, care must be taken to protect the shoulder at all times.

Showering:

- Unless otherwise instructed, you may shower 2 days after surgery, but you must keep the dressing/wounds dry. Do not soak the operative shoulder (no baths, hot-tubs, or pools) until allowed by Dr. McAngus to avoid risk of infection.

Ice Therapy:

- Icing is very important for the first 5-7 postoperative days to decrease swelling and pain. While the post-op dressing is in place, icing can be continuous. Once the dressing is removed on the third operative day, ice can be applied for 15-20 minute periods, 3-4 times per day. Care must be taken with icing to avoid frostbite to the skin.

Sleep:

- Sleeping can be uncomfortable for the first 1-2 weeks after shoulder surgery. It can be helpful to sleep in a recliner-chair or in a semi-upright position.

Diet:

- The anesthetic drugs used during your surgery may cause nausea for the first 24 hours. Following surgery, begin with clear liquids and light foods. You can progress to your normal diet if not nauseated. If nausea and vomiting become severe, or if you show signs of dehydration (lack of urination), please call Dr. McAngus' office.

Post-Operative Appointment:

- Dr. McAngus or Chrisi, LVN will need to reexamine you 14 days after your procedure. Please call the office (numbers on first page) to schedule a follow-up appointment.

Driving:

- No driving until permitted by Dr. McAngus.

Work/School:

- May return to sedentary work or school 3-4 days after surgery, if pain is tolerable

URGENT Information and Contacts:

- A low-grade fever (up to 100.5 degrees) is not uncommon in the first 48 hours. Please call the doctor with any temperature over 101 degrees.
- Please call the doctor if you have severe pain that your pain medication does not relieve, persistent numbness of the leg, fever over 101 degrees, difficulty with the cast or surgical incisions, difficulty breathing, chest pain, excessive nausea/vomiting, or if you experience redness or swelling in your thigh or calf.
- If unexpected problems occur and you need to speak to someone, please call the doctor (254.595.3698). If calling after office hours or on the weekend, you may leave a message, or text the above number. If you do not get a response within 30 minutes, please call the facility/emergency room or present to the nearest emergency room.
- If you have an emergency that requires immediate attention, proceed to the nearest emergency room.
- Follow-Up Care/Questions:
 - Dr. McAngus or Christi, LVN will call you after your surgery. If you have not been contacted within 48 hours of surgery, please reach out to Dr. McAngus' office.
 - If you do not already have a postoperative appointment scheduled, please contact the office during normal office hours. Appointments should be arranged 7-14 days from surgery.
 - Please call (254) 595-3698 with any questions or concerns at any time.


DO NOT REMOVE SPLINT

These exercises are to help with shoulder mobilization



Pendulum Circles:

Shift your body weight in circles to allow your injured arm to swing in circles freely. Perform 3-4 times a day.



Pendulum Forward-Back:

Shift your body weight forward then back to allow your injured arm to swing forward and back freely. Perform 3-4 times a day



Pendulum Side to Side:

Shift your body weight side to side to allow your injured arm to swing side to side freely.